

## VU Research Portal

### Observation of the baryonic B decay $(B)\overline{0} \rightarrow \Lambda(+)(c)(\Lambda)\overline{b}K(-)$

Lees, J.P.; Raven, H.G.; Snoek, H.

**published in**

Physical Review D  
2011

**DOI (link to publisher)**

[10.1103/PhysRevD.84.071102](https://doi.org/10.1103/PhysRevD.84.071102)

**document version**

Publisher's PDF, also known as Version of record

[Link to publication in VU Research Portal](#)

**citation for published version (APA)**

Lees, J. P., Raven, H. G., & Snoek, H. (2011). Observation of the baryonic B decay  $(B)\overline{0} \rightarrow \Lambda(+)(c)(\Lambda)\overline{b}K(-)$ . *Physical Review D*, 84(7), 071102. [071102].  
<https://doi.org/10.1103/PhysRevD.84.071102>

**General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

**Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

**E-mail address:**

[vuresearchportal.ub@vu.nl](mailto:vuresearchportal.ub@vu.nl)

**Observation of the baryonic  $B$  decay  $\bar{B}^0 \rightarrow \Lambda_c^+ \bar{\Lambda} K^-$** 

J. P. Lees,<sup>1</sup> V. Poireau,<sup>1</sup> V. Tisserand,<sup>1</sup> J. Garra Tico,<sup>2</sup> E. Grauges,<sup>2</sup> M. Martinelli,<sup>3a,3b</sup> D. A. Milanes,<sup>3a</sup> A. Palano,<sup>3a,3b</sup> M. Pappagallo,<sup>3a,3b</sup> G. Eigen,<sup>4</sup> B. Stugu,<sup>4</sup> L. Sun,<sup>4</sup> D. N. Brown,<sup>5</sup> L. T. Kerth,<sup>5</sup> Yu. G. Kolomensky,<sup>5</sup> G. Lynch,<sup>5</sup> H. Koch,<sup>6</sup> T. Schroeder,<sup>6</sup> D. J. Asgeirsson,<sup>7</sup> C. Hearty,<sup>7</sup> T. S. Mattison,<sup>7</sup> J. A. McKenna,<sup>7</sup> A. Khan,<sup>8</sup> V. E. Blinov,<sup>9</sup> A. R. Buzykaev,<sup>9</sup> V. P. Druzhinin,<sup>9</sup> V. B. Golubev,<sup>9</sup> E. A. Kravchenko,<sup>9</sup> A. P. Onuchin,<sup>9</sup> S. I. Serednyakov,<sup>9</sup> Yu. I. Skovpen,<sup>9</sup> E. P. Solodov,<sup>9</sup> K. Yu. Todyshev,<sup>9</sup> A. N. Yushkov,<sup>9</sup> M. Bondioli,<sup>10</sup> D. Kirkby,<sup>10</sup> A. J. Lankford,<sup>10</sup> M. Mandelkern,<sup>10</sup> D. P. Stoker,<sup>10</sup> H. Atmacan,<sup>11</sup> J. W. Gary,<sup>11</sup> F. Liu,<sup>11</sup> O. Long,<sup>11</sup> G. M. Vitug,<sup>11</sup> C. Campagnari,<sup>12</sup> T. M. Hong,<sup>12</sup> D. Kovalskyi,<sup>12</sup> J. D. Richman,<sup>12</sup> C. A. West,<sup>12</sup> A. M. Eisner,<sup>12</sup> J. Kroseberg,<sup>13</sup> W. S. Lockman,<sup>13</sup> A. J. Martinez,<sup>13</sup> T. Schalk,<sup>13</sup> B. A. Schumm,<sup>13</sup> A. Seiden,<sup>13</sup> C. H. Cheng,<sup>14</sup> D. A. Doll,<sup>14</sup> B. Echenard,<sup>14</sup> K. T. Flood,<sup>14</sup> D. G. Hitlin,<sup>14</sup> P. Ongmongkolkul,<sup>14</sup> F. C. Porter,<sup>14</sup> A. Y. Rakitin,<sup>14</sup> R. Andreassen,<sup>15</sup> M. S. Dubrovin,<sup>15</sup> Z. Huard,<sup>15</sup> B. T. Meadows,<sup>15</sup> M. D. Sokoloff,<sup>15</sup> P. C. Bloom,<sup>16</sup> W. T. Ford,<sup>16</sup> A. Gaz,<sup>16</sup> M. Nagel,<sup>16</sup> U. Nauenberg,<sup>16</sup> J. G. Smith,<sup>16</sup> S. R. Wagner,<sup>16</sup> R. Ayad,<sup>17,\*</sup> W. H. Toki,<sup>17</sup> B. Spaan,<sup>18</sup> M. J. Kobel,<sup>19</sup> K. R. Schubert,<sup>19</sup> R. Schwierz,<sup>19</sup> D. Bernard,<sup>20</sup> M. Verderi,<sup>20</sup> P. J. Clark,<sup>21</sup> S. Playfer,<sup>21</sup> D. Bettoni,<sup>22a</sup> C. Bozzi,<sup>22a</sup> R. Calabrese,<sup>22a,22b</sup> G. Cibinetto,<sup>22a,22b</sup> E. Fioravanti,<sup>22a,22b</sup> I. Garzia,<sup>22a,22b</sup> E. Luppi,<sup>22a,22b</sup> M. Munerato,<sup>22a,22b</sup> M. Negrini,<sup>22a,22b</sup> L. Piemontese,<sup>22a</sup> R. Baldini-Ferroli,<sup>23</sup> A. Calcaterra,<sup>23</sup> R. de Sangro,<sup>23</sup> G. Finocchiaro,<sup>23</sup> M. Nicolaci,<sup>23</sup> P. Patteri,<sup>23</sup> I. M. Peruzzi,<sup>23,†</sup> M. Piccolo,<sup>23</sup> M. Rama,<sup>23</sup> A. Zallo,<sup>23</sup> R. Contri,<sup>24a,24b</sup> E. Guido,<sup>24a,24b</sup> M. Lo Vetere,<sup>24a,24b</sup> M. R. Monge,<sup>24a,24b</sup> S. Passaggio,<sup>24a</sup> C. Patrignani,<sup>24a,24b</sup> E. Robutti,<sup>24a</sup> B. Bhuyan,<sup>25</sup> V. Prasad,<sup>25</sup> C. L. Lee,<sup>26</sup> M. Morii,<sup>26</sup> A. J. Edwards,<sup>27</sup> A. Adametz,<sup>28</sup> J. Marks,<sup>28</sup> U. Uwer,<sup>28</sup> F. U. Bernlochner,<sup>29</sup> M. Ebert,<sup>29</sup> H. M. Lacker,<sup>29</sup> T. Lueck,<sup>29</sup> P. D. Dauncey,<sup>30</sup> M. Tibbetts,<sup>30</sup> P. K. Behera,<sup>31</sup> U. Mallik,<sup>31</sup> C. Chen,<sup>32</sup> J. Cochran,<sup>32</sup> W. T. Meyer,<sup>32</sup> S. Prell,<sup>32</sup> E. I. Rosenberg,<sup>32</sup> A. E. Rubin,<sup>32</sup> A. V. Gritsan,<sup>33</sup> Z. J. Guo,<sup>33</sup> N. Arnaud,<sup>34</sup> M. Davier,<sup>34</sup> G. Grosdidier,<sup>34</sup> F. Le Diberder,<sup>34</sup> A. M. Lutz,<sup>34</sup> B. Malaescu,<sup>34</sup> P. Roudeau,<sup>34</sup> M. H. Schune,<sup>34</sup> A. Stocchi,<sup>34</sup> G. Wormser,<sup>34</sup> D. J. Lange,<sup>35</sup> D. M. Wright,<sup>35</sup> I. Bingham,<sup>36</sup> C. A. Chavez,<sup>36</sup> J. P. Coleman,<sup>36</sup> J. R. Fry,<sup>36</sup> E. Gabathuler,<sup>36</sup> D. E. Hutchcroft,<sup>36</sup> D. J. Payne,<sup>36</sup> C. Touramanis,<sup>36</sup> A. J. Bevan,<sup>37</sup> F. Di Lodovico,<sup>37</sup> R. Sacco,<sup>37</sup> M. Sigamani,<sup>37</sup> G. Cowan,<sup>37</sup> S. Paramesvaran,<sup>38</sup> D. N. Brown,<sup>39</sup> C. L. Davis,<sup>39</sup> A. G. Denig,<sup>40</sup> M. Fritsch,<sup>40</sup> W. Gradl,<sup>40</sup> A. Hafner,<sup>40</sup> E. Prencipe,<sup>40</sup> K. E. Alwyn,<sup>41</sup> D. Bailey,<sup>41</sup> R. J. Barlow,<sup>41,‡</sup> G. Jackson,<sup>41</sup> G. D. Lafferty,<sup>41</sup> R. Cenci,<sup>42</sup> B. Hamilton,<sup>42</sup> A. Jawahery,<sup>42</sup> D. A. Roberts,<sup>42</sup> G. Simi,<sup>42</sup> C. Dallapiccola,<sup>43</sup> R. Cowan,<sup>44</sup> D. Dujmic,<sup>44</sup> G. Sciolla,<sup>44</sup> D. Lindemann,<sup>45</sup> P. M. Patel,<sup>45</sup> S. H. Robertson,<sup>45</sup> M. Schram,<sup>45</sup> P. Biassoni,<sup>46a,46b</sup> A. Lazzaro,<sup>46a,46b</sup> V. Lombardo,<sup>46a</sup> N. Neri,<sup>46a,46b</sup> F. Palombo,<sup>46a,46b</sup> S. Stracka,<sup>46a,46b</sup> L. Cremaldi,<sup>47</sup> R. Godang,<sup>47,§</sup> R. Kroeger,<sup>47</sup> P. Sonnek,<sup>47</sup> D. J. Summers,<sup>47</sup> X. Nguyen,<sup>48</sup> P. Taras,<sup>48</sup> G. De Nardo,<sup>49a,49b</sup> D. Monorchio,<sup>49a,49b</sup> G. Onorato,<sup>49a,49b</sup> C. Sciacca,<sup>49a,49b</sup> G. Raven,<sup>50</sup> H. L. Snoek,<sup>50</sup> C. P. Jessop,<sup>51</sup> K. J. Knoepfel,<sup>51</sup> J. M. LoSecco,<sup>51</sup> W. F. Wang,<sup>51</sup> K. Honscheid,<sup>52</sup> R. Kass,<sup>52</sup> J. Brau,<sup>53</sup> R. Frey,<sup>53</sup> N. B. Sinev,<sup>53</sup> D. Strom,<sup>53</sup> E. Torrence,<sup>53</sup> E. Feltresi,<sup>54a,54b</sup> N. Gagliardi,<sup>54a,54b</sup> M. Margoni,<sup>54a,54b</sup> M. Morandin,<sup>54a</sup> M. Posocco,<sup>54a</sup> M. Rotondo,<sup>54a</sup> F. Simonetto,<sup>54a,54b</sup> R. Stroili,<sup>55</sup> E. Ben-Haim,<sup>55</sup> M. Bomben,<sup>55</sup> G. R. Bonneaud,<sup>55</sup> H. Briand,<sup>55</sup> G. Calderini,<sup>55</sup> J. Chauveau,<sup>55</sup> O. Hamon,<sup>55</sup> Ph. Leruste,<sup>55</sup> G. Marchiori,<sup>55</sup> J. Ocariz,<sup>55</sup> S. Sitt,<sup>55</sup> M. Biasini,<sup>56a,56b</sup> E. Manoni,<sup>56a,56b</sup> S. Pacetti,<sup>56a,56b</sup> A. Rossi,<sup>56a,56b</sup> C. Angelini,<sup>57a,57b</sup> G. Batignani,<sup>57a,57b</sup> S. Bettarini,<sup>57a,57b</sup> M. Carpinelli,<sup>57a,57b,||</sup> G. Casarosa,<sup>57a,57b</sup> A. Cervelli,<sup>57a,57b</sup> F. Forti,<sup>57a,57b</sup> M. A. Giorgi,<sup>57a,57b</sup> A. Lusiani,<sup>57a,57c</sup> B. Oberhof,<sup>57a,57b</sup> E. Paoloni,<sup>57a,57b</sup> A. Perez,<sup>57a</sup> G. Rizzo,<sup>57a,57b</sup> J. J. Walsh,<sup>57a</sup> D. Lopes Pegna,<sup>57a,57b</sup> C. Lu,<sup>58</sup> J. Olsen,<sup>58</sup> A. J. S. Smith,<sup>58</sup> A. V. Telnov,<sup>58</sup> F. Anulli,<sup>59a</sup> G. Cavoto,<sup>59a</sup> R. Faccini,<sup>59a</sup> F. Ferrarotto,<sup>59a,59b</sup> F. Ferroni,<sup>59a</sup> M. Gaspero,<sup>59a,59b</sup> L. Li Gioi,<sup>59a</sup> M. A. Mazzoni,<sup>59a</sup> G. Piredda,<sup>59a</sup> C. Büniger,<sup>60</sup> O. Grünberg,<sup>60</sup> T. Hartmann,<sup>60</sup> T. Leddig,<sup>60</sup> H. Schröder,<sup>60</sup> R. Waldi,<sup>60</sup> T. Adye,<sup>61</sup> E. O. Olaiya,<sup>61</sup> F. F. Wilson,<sup>61</sup> S. Emery,<sup>62</sup> G. Hamel de Monchenault,<sup>62</sup> G. Vasseur,<sup>62</sup> Ch. Yèche,<sup>62</sup> D. Aston,<sup>63</sup> D. J. Bard,<sup>63</sup> R. Bartoldus,<sup>63</sup> C. Cartaro,<sup>63</sup> M. R. Convery,<sup>63</sup> J. Dorfan,<sup>63</sup> G. P. Dubois-Felsmann,<sup>63</sup> W. Dunwoodie,<sup>63</sup> R. C. Field,<sup>63</sup> M. Franco Sevilla,<sup>63</sup> B. G. Fulsom,<sup>63</sup> A. M. Gabareen,<sup>63</sup> M. T. Graham,<sup>63</sup> P. Grenier,<sup>63</sup> C. Hast,<sup>63</sup> W. R. Innes,<sup>63</sup> M. H. Kelsey,<sup>63</sup> H. Kim,<sup>63</sup> P. Kim,<sup>63</sup> M. L. Kocian,<sup>63</sup> D. W. G. S. Leith,<sup>63</sup> P. Lewis,<sup>63</sup> S. Li,<sup>63</sup> B. Lindquist,<sup>63</sup> S. Luitz,<sup>63</sup> V. Luth,<sup>63</sup> H. L. Lynch,<sup>63</sup> D. B. MacFarlane,<sup>63</sup> D. R. Muller,<sup>63</sup> H. Neal,<sup>63</sup> S. Nelson,<sup>63</sup> I. Ofte,<sup>63</sup> M. Perl,<sup>63</sup> T. Pulliam,<sup>63</sup> B. N. Ratcliff,<sup>63</sup> A. Roodman,<sup>63</sup> A. A. Salnikov,<sup>63</sup> V. Santoro,<sup>63</sup> R. H. Schindler,<sup>63</sup> A. Snyder,<sup>63</sup> D. Su,<sup>63</sup> M. K. Sullivan,<sup>63</sup> J. Va'vra,<sup>63</sup> A. P. Wagner,<sup>63</sup> M. Weaver,<sup>63</sup> W. J. Wisniewski,<sup>63</sup> M. Wittgen,<sup>63</sup> D. H. Wright,<sup>63</sup> H. W. Wulsin,<sup>63</sup> A. K. Yarritu,<sup>63</sup> C. C. Young,<sup>63</sup> V. Ziegler,<sup>63</sup> W. Park,<sup>64</sup> M. V. Purohit,<sup>64</sup> R. M. White,<sup>64</sup> J. R. Wilson,<sup>64</sup> A. Randle-Conde,<sup>65</sup> S. J. Sekula,<sup>65</sup> M. Bellis,<sup>66</sup> J. F. Benitez,<sup>66</sup> P. R. Burchat,<sup>66</sup> T. S. Miyashita,<sup>66</sup> M. S. Alam,<sup>67</sup> J. A. Ernst,<sup>67</sup> R. Gorodeisky,<sup>68</sup> N. Guttman,<sup>68</sup> D. R. Peimer,<sup>68</sup> A. Soffer,<sup>68</sup> P. Lund,<sup>69</sup> S. M. Spanier,<sup>69</sup> R. Eckmann,<sup>70</sup> J. L. Ritchie,<sup>70</sup> A. M. Ruland,<sup>70</sup> C. J. Schilling,<sup>70</sup> R. F. Schwitters,<sup>70</sup> B. C. Wray,<sup>70</sup> J. M. Izen,<sup>71</sup> X. C. Lou,<sup>71</sup> F. Bianchi,<sup>72a,72b</sup> D. Gamba,<sup>72a,72b</sup> L. Lanceri,<sup>73a,73b</sup> L. Vitale,<sup>73a,73b</sup> F. Martinez-Vidal,<sup>74</sup> A. Oyanguren,<sup>74</sup> H. Ahmed,<sup>75</sup> J. Albert,<sup>75</sup> Sw. Banerjee,<sup>75</sup> H. H. F. Choi,<sup>75</sup>

G. J. King,<sup>75</sup> R. Kowalewski,<sup>75</sup> M. J. Lewczuk,<sup>75</sup> C. Lindsay,<sup>75</sup> I. M. Nugent,<sup>75</sup> J. M. Roney,<sup>75</sup> R. J. Sobie,<sup>75</sup>  
 T. J. Gershon,<sup>75</sup> P. F. Harrison,<sup>76</sup> T. E. Latham,<sup>76</sup> E. M. T. Puccio,<sup>76</sup> H. R. Band,<sup>77</sup> S. Dasu,<sup>77</sup> Y. Pan,<sup>77</sup> R. Prepost,<sup>77</sup>  
 C. O. Vuosalo,<sup>77</sup> and S. L. Wu<sup>77</sup>

(BABAR Collaboration)

- <sup>1</sup>Laboratoire d'Annecy-le-Vieux de Physique des Particules (LAPP), Université de Savoie, CNRS/IN2P3, F-74941 Annecy-Le-Vieux, France  
<sup>2</sup>Universitat de Barcelona, Facultat de Física, Departament ECM, E-08028 Barcelona, Spain  
<sup>3a</sup>INFN Sezione di Bari, I-70126 Bari, Italy  
<sup>3b</sup>Dipartimento di Fisica, Università di Bari, I-70126 Bari, Italy  
<sup>4</sup>University of Bergen, Institute of Physics, N-5007 Bergen, Norway  
<sup>5</sup>Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720, USA  
<sup>6</sup>Ruhr Universität Bochum, Institut für Experimentalphysik 1, D-44780 Bochum, Germany  
<sup>7</sup>University of British Columbia, Vancouver, British Columbia, Canada V6T 1Z1  
<sup>8</sup>Brunel University, Uxbridge, Middlesex UB8 3PH, United Kingdom  
<sup>9</sup>Budker Institute of Nuclear Physics, Novosibirsk 630090, Russia  
<sup>10</sup>University of California at Irvine, Irvine, California 92697, USA  
<sup>11</sup>University of California at Riverside, Riverside, California 92521, USA  
<sup>12</sup>University of California at Santa Barbara, Santa Barbara, California 93106, USA  
<sup>13</sup>University of California at Santa Cruz, Institute for Particle Physics, Santa Cruz, California 95064, USA  
<sup>14</sup>California Institute of Technology, Pasadena, California 91125, USA  
<sup>15</sup>University of Cincinnati, Cincinnati, Ohio 45221, USA  
<sup>16</sup>University of Colorado, Boulder, Colorado 80309, USA  
<sup>17</sup>Colorado State University, Fort Collins, Colorado 80523, USA  
<sup>18</sup>Technische Universität Dortmund, Fakultät Physik, D-44221 Dortmund, Germany  
<sup>19</sup>Technische Universität Dresden, Institut für Kern- und Teilchenphysik, D-01062 Dresden, Germany  
<sup>20</sup>Laboratoire Leprince-Ringuet, Ecole Polytechnique, CNRS/IN2P3, F-91128 Palaiseau, France  
<sup>21</sup>University of Edinburgh, Edinburgh EH9 3JZ, United Kingdom  
<sup>22a</sup>INFN Sezione di Ferrara, I-44100 Ferrara, Italy  
<sup>22b</sup>Dipartimento di Fisica, Università di Ferrara, I-44100 Ferrara, Italy  
<sup>23</sup>INFN Laboratori Nazionali di Frascati, I-00044 Frascati, Italy  
<sup>24a</sup>INFN Sezione di Genova, I-16146 Genova, Italy  
<sup>24b</sup>Dipartimento di Fisica, Università di Genova, I-16146 Genova, Italy  
<sup>25</sup>Indian Institute of Technology Guwahati, Guwahati, Assam, 781 039, India  
<sup>26</sup>Harvard University, Cambridge, Massachusetts 02138, USA  
<sup>27</sup>Harvey Mudd College, Claremont, California 91711, USA  
<sup>28</sup>Universität Heidelberg, Physikalisches Institut, Philosophenweg 12, D-69120 Heidelberg, Germany  
<sup>29</sup>Humboldt-Universität zu Berlin, Institut für Physik, Newtonstrasse 15, D-12489 Berlin, Germany  
<sup>30</sup>Imperial College London, London, SW7 2AZ, United Kingdom  
<sup>31</sup>University of Iowa, Iowa City, Iowa 52242, USA  
<sup>32</sup>Iowa State University, Ames, Iowa 50011-3160, USA  
<sup>33</sup>Johns Hopkins University, Baltimore, Maryland 21218, USA  
<sup>34</sup>Laboratoire de l'Accélérateur Linéaire, IN2P3/CNRS et Université Paris-Sud 11, Centre Scientifique d'Orsay, B.P. 34, F-91898 Orsay Cedex, France  
<sup>35</sup>Lawrence Livermore National Laboratory, Livermore, California 94550, USA  
<sup>36</sup>University of Liverpool, Liverpool L69 7ZE, United Kingdom  
<sup>37</sup>Queen Mary, University of London, London, E1 4NS, United Kingdom  
<sup>38</sup>University of London, Royal Holloway and Bedford New College, Egham, Surrey TW20 0EX, United Kingdom  
<sup>39</sup>University of Louisville, Louisville, Kentucky 40292, USA  
<sup>40</sup>Johannes Gutenberg-Universität Mainz, Institut für Kernphysik, D-55099 Mainz, Germany  
<sup>41</sup>University of Manchester, Manchester M13 9PL, United Kingdom  
<sup>42</sup>University of Maryland, College Park, Maryland 20742, USA  
<sup>43</sup>University of Massachusetts, Amherst, Massachusetts 01003, USA  
<sup>44</sup>Massachusetts Institute of Technology, Laboratory for Nuclear Science, Cambridge, Massachusetts 02139, USA  
<sup>45</sup>McGill University, Montréal, Québec, Canada H3A 2T8  
<sup>46a</sup>INFN Sezione di Milano, I-20133 Milano, Italy  
<sup>46b</sup>Dipartimento di Fisica, Università di Milano, I-20133 Milano, Italy  
<sup>47</sup>University of Mississippi, University, Mississippi 38677, USA  
<sup>48</sup>Université de Montréal, Physique des Particules, Montréal, Québec, Canada H3C 3J7

- <sup>49a</sup>INFN Sezione di Napoli, I-80126 Napoli, Italy  
<sup>49b</sup>Dipartimento di Scienze Fisiche, Università di Napoli Federico II, I-80126 Napoli, Italy  
<sup>50</sup>NIKHEF, National Institute for Nuclear Physics and High Energy Physics, NL-1009 DB Amsterdam, The Netherlands  
<sup>51</sup>University of Notre Dame, Notre Dame, Indiana 46556, USA  
<sup>52</sup>Ohio State University, Columbus, Ohio 43210, USA  
<sup>53</sup>University of Oregon, Eugene, Oregon 97403, USA  
<sup>54a</sup>INFN Sezione di Padova, I-35131 Padova, Italy  
<sup>54b</sup>Dipartimento di Fisica, Università di Padova, I-35131 Padova, Italy  
<sup>55</sup>Laboratoire de Physique Nucléaire et de Hautes Energies, IN2P3/CNRS, Université Pierre et Marie Curie-Paris6, Université Denis Diderot-Paris7, F-75252 Paris, France  
<sup>56a</sup>INFN Sezione di Perugia, I-06100 Perugia, Italy  
<sup>56b</sup>Dipartimento di Fisica, Università di Perugia, I-06100 Perugia, Italy  
<sup>57a</sup>INFN Sezione di Pisa, I-56127 Pisa, Italy  
<sup>57b</sup>Dipartimento di Fisica, I-56127 Pisa, Italy  
<sup>57c</sup>Università di Pisa, Scuola Normale Superiore di Pisa, I-56127 Pisa, Italy  
<sup>58</sup>Princeton University, Princeton, New Jersey 08544, USA  
<sup>59a</sup>INFN Sezione di Roma, I-00185 Roma, Italy  
<sup>59b</sup>Dipartimento di Fisica, Università di Roma La Sapienza, I-00185 Roma, Italy  
<sup>60</sup>Universität Rostock, D-18051 Rostock, Germany  
<sup>61</sup>Rutherford Appleton Laboratory, Chilton, Didcot, Oxon, OX11 0QX, United Kingdom  
<sup>62</sup>CEA, Ifu, SPP, Centre de Saclay, F-91191 Gif-sur-Yvette, France  
<sup>63</sup>SLAC National Accelerator Laboratory, Stanford, California 94309 USA  
<sup>64</sup>University of South Carolina, Columbia, South Carolina 29208, USA  
<sup>65</sup>Southern Methodist University, Dallas, Texas 75275, USA  
<sup>66</sup>Stanford University, Stanford, California 94305-4060, USA  
<sup>67</sup>State University of New York, Albany, New York 12222, USA  
<sup>68</sup>Tel Aviv University, School of Physics and Astronomy, Tel Aviv, 69978, Israel  
<sup>69</sup>University of Tennessee, Knoxville, Tennessee 37996, USA  
<sup>70</sup>University of Texas at Austin, Austin, Texas 78712, USA  
<sup>71</sup>University of Texas at Dallas, Richardson, Texas 75083, USA  
<sup>72a</sup>INFN Sezione di Torino, I-10125 Torino, Italy  
<sup>72b</sup>Dipartimento di Fisica Sperimentale, Università di Torino, I-10125 Torino, Italy  
<sup>73a</sup>INFN Sezione di Trieste, I-34127 Trieste, Italy  
<sup>73b</sup>Dipartimento di Fisica, Università di Trieste, I-34127 Trieste, Italy  
<sup>74</sup>IFIC, Universitat de Valencia-CSIC, E-46071 Valencia, Spain  
<sup>75</sup>University of Victoria, Victoria, British Columbia, Canada V8W 3P6  
<sup>76</sup>Department of Physics, University of Warwick, Coventry CV4 7AL, United Kingdom  
<sup>77</sup>University of Wisconsin, Madison, Wisconsin 53706, USA

(Received 17 August 2011; published 11 October 2011; publisher error corrected 7 February 2012)

We report the observation of the baryonic  $B$  decay  $\bar{B}^0 \rightarrow \Lambda_c^+ \bar{\Lambda} K^-$  with a significance larger than 7 standard deviations based on  $471 \times 10^6$   $B\bar{B}$  pairs collected with the BABAR detector at the PEP-II storage ring at SLAC. We measure the branching fraction for the decay  $\bar{B}^0 \rightarrow \Lambda_c^+ \bar{\Lambda} K^-$  to be  $(3.8 \pm 0.8_{\text{stat}} \pm 0.2_{\text{sys}} \pm 1.0_{\Lambda_c^+}) \times 10^{-5}$ . The uncertainties are statistical, systematic, and due to the uncertainty in the  $\Lambda_c^+$  branching fraction. We find that the  $\Lambda_c^+ K^-$  invariant-mass distribution shows an enhancement above 3.5 GeV/ $c^2$ .

DOI: 10.1103/PhysRevD.84.071102

PACS numbers: 13.25.Hw, 13.60.Rj, 14.20.Lq

While baryons are produced in  $(6.8 \pm 0.6)\%$  [1] of all  $B$ -meson decays, little is known about the detailed mechanics of these decays and more generally about hadron

fragmentation into baryons. We can increase our understanding of baryon production in  $B$  decays by comparing decay rates for related exclusive final states. In this paper we present a measurement of the decay  $\bar{B}^0 \rightarrow \Lambda_c^+ \bar{\Lambda} K^-$  [2]. Currently, no experimental results are available for this decay.

This analysis is based on a data set of about  $429 \text{ fb}^{-1}$ , corresponding to  $471 \times 10^6$   $B\bar{B}$  pairs, collected with the BABAR detector at the PEP-II asymmetric-energy  $e^+e^-$  storage ring, operated at a center-of-mass energy equal to the  $Y(4S)$  mass. The signal efficiency is determined with a

\*Now at Temple University, Philadelphia, PA 19122, USA.

†Also with Università di Perugia, Dipartimento di Fisica, Perugia, Italy.

‡Now at the University of Huddersfield, Huddersfield HD1 3DH, United Kingdom.

§Now at University of South Alabama, Mobile, AL 36688, USA.

||Also with Università di Sassari, Sassari, Italy.



Monte Carlo simulation based on EVTGEN [3] for the event generation and GEANT4 [4] for the detector simulation. The  $\bar{B}^0 \rightarrow \Lambda_c^+ \bar{\Lambda} K^-$  Monte Carlo events are generated uniformly in the  $\Lambda_c^+ \bar{\Lambda} K^-$  phase space. Monte Carlo simulated events are used to study background contributions as well.

The *BABAR* detector is described in detail elsewhere [5]. Charged particle trajectories are measured by a five-layer double-sided silicon vertex tracker and a 40-layer drift chamber, both immersed in a 1.5 T axial magnetic field. Charged particle identification is provided by ionization energy measurements along with Cherenkov radiation detection by an internally reflecting ring-imaging detector.

The  $\Lambda_c^+$  is reconstructed in the decay mode  $\Lambda_c^+ \rightarrow p K^- \pi^+$  and the  $\bar{\Lambda}$  in the decay mode  $\bar{\Lambda} \rightarrow \bar{p} \pi^+$ . For the identification of proton, kaon, and pion candidates, we use selection criteria based on the measurements of the specific ionization in the tracking detectors and of the Cherenkov radiation in the internally reflecting ring-imaging detector [6].

For the identification of the  $p$  coming from the  $\Lambda_c^+$  the average efficiency is about 95%, while the probability of misidentifying a kaon as a proton is less than 2%. The average efficiency for the  $K^-$  identification is about 90%. The probability of misidentifying a pion as a kaon is about 5%. These are the dominant misidentification probabilities for each particle type. The  $\Lambda_c^+$  daughters and the  $\bar{\Lambda}$  daughters are each fit to a common vertex, and the  $\Lambda_c^+$  and the  $\bar{\Lambda}$  candidate invariant mass is required to lie within  $3\sigma$  of the world average mass [1], i.e., in the range 2.273–2.299 GeV/ $c^2$  and 1.113–1.119 GeV/ $c^2$ , respectively. For the reconstruction of the  $B$  candidate, the mass of the  $\Lambda_c^+$  candidate is constrained to its nominal value [1] and is combined with a  $\bar{\Lambda}$  and a  $K^-$  candidate. Since the  $\bar{\Lambda}$  candidate mass is already well measured, it is not constrained.

The  $\Lambda_c^+$ ,  $\bar{\Lambda}$  and  $K^-$  candidates are then fitted to a common vertex, and the confidence level of this fit is required to exceed 0.2%.

A possible source for fake signal events is the decay  $\bar{B}^0 \rightarrow \Lambda_c^+ \bar{p} K^- \pi^+$  [7], which has the same final state as the decay under investigation. In order to suppress this background we require that the distance between the  $B$  vertex and the  $\bar{\Lambda}$  vertex in the  $xy$  plane (with  $z$  parallel to the beam axis) exceeds 0.4 cm. This constraint reduces combinatoric background by 18% and the background from  $\bar{B}^0 \rightarrow \Lambda_c^+ \bar{p} K^- \pi^+$  by 99.6%. The expected remaining background from this decay is determined to be  $0.1 \pm 0.1$  events [7].

The separation of signal and background in the candidate sample is obtained by using two kinematic variables:

$\Delta E = E_B^* - \sqrt{s}/2$  and  $m_{\text{ES}} = \sqrt{(s/2 + \mathbf{p}_i \cdot \mathbf{p}_B)^2/E_i^2 - |\mathbf{p}_B|^2}$ , where  $\sqrt{s}$  is the  $e^+e^-$  center-of-mass energy and  $E_B^*$  the energy of the  $B$  candidate in the center-of-mass system.  $(E_i, \mathbf{p}_i)$  is the four-momentum vector of the  $e^+e^-$  system and  $\mathbf{p}_B$  the  $B$ -candidate momentum vector, both measured




FIG. 1 (color online). The  $\Delta E$  distribution for  $\Lambda_c^+ \bar{\Lambda} K^-$  candidates in the data with all selection criteria applied (points). The solid line is the overall fit result, while the dashed line is the background component.

in the laboratory frame. For true  $B$  decays  $m_{\text{ES}}$  is centered at the  $B$ -meson mass, and  $\Delta E$  is centered at zero.  $B$  candidates are required to have an  $m_{\text{ES}}$  value between 5.272 and 5.288 GeV/ $c^2$ .

Figure 1 shows the  $\Delta E$  distribution of the selected candidates, fitted in the range from  $-0.12$  to  $0.30$  GeV. We fit the signal with a Gaussian with the mean  $\mu$  and width  $\sigma$  fixed to the values obtained from a fit to the Monte Carlo simulation ( $\mu = 0.247$  MeV and  $\sigma = 8.381$  MeV), leaving only the signal yield floating. The background is described by a first-order polynomial. A binned maximum likelihood fit with this probability density function gives a signal yield of  $51 \pm 9$  events. (For the branching fraction measurement described later, we use the excess number of candidates above background as the estimate of the number of signal events.) The confidence level for the null hypothesis, considering statistical uncertainties only, is  $2.6 \times 10^{-15}$ , which corresponds to a statistical significance of 8 standard deviations. A possible background from  $\bar{B}^0 \rightarrow \Lambda_c^+ \bar{\Sigma}^0 K^-$ , which rises slowly up to  $\Delta E \approx -0.06$  GeV and drops sharply between  $\Delta E = -0.05$  and  $-0.02$  GeV, is not visible in Fig. 1.

Since the decay dynamics of baryonic  $B$  decays are largely unknown, we investigate the invariant-mass distribution of the two-body systems. Intermediate states would appear as differences in the invariant-mass distribution for data and for the  $\bar{B}^0 \rightarrow \Lambda_c^+ \bar{\Lambda} K^-$  Monte Carlo simulation, in which the final state is generated according to three-body phase space. Using the same function that we used in Fig. 1, we fit the  $\Delta E$  distributions for ten ranges of the three two-body masses. The results are compared to the phase-space Monte Carlo simulation in Fig. 2. While the  $m(\Lambda_c^+ \bar{\Lambda})$  and  $m(\bar{\Lambda} K^-)$  distributions show no significant deviations, the  $m(\Lambda_c^+ K^-)$  distribution shows the data concentrated in the upper half of the allowed mass range, contrary to the Monte Carlo simulation. A possible explanation for this is a


OBSERVATION OF THE BARYONIC  $B$  DECAY ...

FIG. 2 (color online). The  $m(\Lambda_c^+ \bar{\Lambda})$ ,  $m(\bar{\Lambda} K^-)$ , and  $m(\Lambda_c^+ K^-)$  distributions in the data (points) in comparison with the Monte Carlo sample (red histogram).

resonant decay via a baryon resonance that has not yet been observed. Another possibility is enhanced rates at both  $m(\Lambda_c^+ \bar{\Lambda})$  and  $m(\bar{\Lambda} K^-)$  thresholds.

Because the efficiency varies over the Dalitz plot and the distribution of candidates in data is unknown *a priori*, we must use the distribution of the data events in the Dalitz plot to estimate the efficiency. The small number of candidates, combined with resolution and edge effects, makes the simple weighting of events by the inverse of

PHYSICAL REVIEW D **84**, 071102(R) (2011)

TABLE I. Two-body invariant-mass distributions used for the weighting and the corresponding reconstruction efficiency  $\varepsilon$ . The dots indicate how often the respective mass distributions are used to determine the weights.

$m(\Lambda_c^+ K^-)$	$m(\Lambda_c^+ \bar{\Lambda})$	$m(\bar{\Lambda} K^-)$	Efficiency $\varepsilon$ in %
			10.90
•			8.60
•	•		9.21
•	•	•	9.19
••	•	•	8.81
••	••	•	8.80
••	••	••	8.77

the efficiency problematic. Instead, we determine a set of weights to apply to the simulated events so that the resulting weighted Monte Carlo distributions mimic the data. We make the assumption that the dependence of the decay dynamics on the two-body invariant masses can be factorized into the product of three functions that each depend on one invariant mass and weight the Monte Carlo events with the function:

$$w[m(\Lambda_c^+ K^-), m(\Lambda_c^+ \bar{\Lambda}), m(\bar{\Lambda} K^-)] = w_a[m(\Lambda_c^+ K^-)] \cdot w_b[m(\Lambda_c^+ \bar{\Lambda})] \cdot w_c[m(\bar{\Lambda} K^-)]. \quad (1)$$

By dividing the background-subtracted  $m(\Lambda_c^+ K^-)$  distribution (Fig. 2) by the corresponding distribution from the phase-space Monte Carlo simulation, we obtain the weights  $w_a[m(\Lambda_c^+ K^-)]$ , which are used to weight the Monte Carlo candidates. (If a negative weight is required, the weight is constrained to zero.) Next, we use the weighted Monte Carlo candidates to determine  $w_b[m(\Lambda_c^+ \bar{\Lambda})]$  in the same way and then use  $w_a[m(\Lambda_c^+ K^-)] \cdot w_b[m(\Lambda_c^+ \bar{\Lambda})]$  to determine  $w_c[m(\bar{\Lambda} K^-)]$ . After each weighting we determine the reconstruction efficiency by a fit to the  $\Delta E$  distribution for weighted Monte Carlo. Starting with these weights, the weighting is repeated until the reconstruction efficiency converges and the two-body mass distributions in the data and Monte Carlo agree within statistical uncertainties. The efficiency after each weighting is shown in Table I. Since the  $m(\Lambda_c^+ K^-)$  distribution in the data shows the strongest deviations compared to the phase-space Monte Carlo simulation, we use the efficiency  $\varepsilon$  obtained after the second weighting in  $m(\Lambda_c^+ K^-)$  [ $\varepsilon = 8.81\%$ ]. The comparison between the data and weighted Monte Carlo events in the two-body masses can be seen in Fig. 3. Note that, by construction, the data and simulation agree exactly for the  $m(\Lambda_c^+ K^-)$  distribution. The close agreement in the other two distributions shows that the form given in Eq. (1) is adequate to describe any correlations between variables in the data. The effect of the statistical uncertainties in the data on the efficiency determination is described below.

For the branching fraction calculation we determine the number of reconstructed events by a fit to  $\Delta E$  with a




FIG. 3 (color online). The background-subtracted  $m(\Lambda_c^+ \bar{\Lambda})$ ,  $m(\Lambda_c^+ K^-)$ , and  $m(\Lambda_c^+ \bar{K})$  distributions for the data (points) and for the final weighted Monte Carlo sample (red histogram).

first-order polynomial for the background. To avoid a potential bias introduced by an assumption of the signal shape, we fit the region  $-0.12 < \Delta E < 0.30$  GeV exclusive of the signal region  $-0.03 < \Delta E < 0.03$  GeV. By extrapolating the background yield into the signal region and subtracting it from the integral of the histogram in this region, we obtain a signal yield of  $N_{\text{sig}} = 50 \pm 11$ . This results in a branching fraction of

TABLE II. Summary of the relative systematic uncertainties on the branching fraction  $\mathcal{B}(\bar{B}^0 \rightarrow \Lambda_c^+ \bar{\Lambda} K^-)$ .

Source	Relative syst. uncert.
Track reconstruction	0.9%
Charged particle ID	2.4%
$N_{B\bar{B}}$	0.6%
$\mathcal{B}(\bar{\Lambda} \rightarrow \bar{p} \pi^+)$	0.8%
Monte Carlo statistics	0.5%
$\bar{B}^0 \rightarrow \Lambda_c^+ \bar{\Sigma}^0 K^-$	1.0%
$B \rightarrow \Lambda_c^+ \bar{\Lambda} K^- \pi$	1.8%
$\Delta E$ background description	2.0%
$\mu(\Delta E)$ shift	0.8%
Efficiency determination	0.5%
Total	4.1%

$$\begin{aligned}
 \mathcal{B}(\bar{B}^0 \rightarrow \Lambda_c^+ \bar{\Lambda} K^-) &= \frac{N_{\text{sig}}/\varepsilon}{N_{B\bar{B}} \cdot \mathcal{B}(\Lambda_c^+ \rightarrow p K^- \pi^+) \cdot \mathcal{B}(\bar{\Lambda} \rightarrow \bar{p} \pi^+)} \\
 &= (3.8 \pm 0.8_{\text{stat}} \pm 1.0_{\Lambda_c^+}) \times 10^{-5}, \quad (2)
 \end{aligned}$$

with  $N_{B\bar{B}} = N_{\bar{B}^0} + N_{B^0} = (471 \pm 3) \times 10^6$ , assuming equal production of  $B^0 \bar{B}^0$  and  $B^+ B^-$  in the decay of the  $\Upsilon(4S)$ . The branching fractions  $\mathcal{B}(\Lambda_c^+ \rightarrow p K^- \pi^+) = (5.0 \pm 1.3)\%$  and  $\mathcal{B}(\bar{\Lambda} \rightarrow \bar{p} \pi^+) = (63.9 \pm 0.5)\%$  are the world averages from Ref. [1].

Several sources of systematic uncertainties are investigated and summarized in Table II. Most of the uncertainties are derived from comparisons between Monte Carlo simulations and control samples in the data. Systematic uncertainties arise from uncertainties in charged particle reconstruction efficiencies (0.9%) and charged particle identification efficiencies (2.4%) and from statistical uncertainties in the Monte Carlo simulation (0.5%). The systematic uncertainty on the number of  $B\bar{B}$  pairs is 0.6%. The systematic uncertainty from the  $\bar{\Lambda}$  branching fraction amounts to 0.8%.

The systematic uncertainty introduced by neglecting a possible  $\bar{B}^0 \rightarrow \Lambda_c^+ \bar{\Sigma}^0 K^-$  background is determined by adding a probability density function for this background to the fit function used for the  $\Delta E$  fit shown in Fig. 1. Allowing non-negative contributions from this background, the fit returns a value of  $0.0_{-0.0}^{+1.8}$ . For a conservative limit on this systematic uncertainty we fix the yield to 1.8 and take the change in the number of signal events as a systematic uncertainty (1.0%). The  $\Delta E$  distribution in Fig. 1 shows an enhancement below  $-0.14$  GeV, caused by decays of the type  $B \rightarrow \Lambda_c^+ \bar{\Lambda} K^- \pi$ . Because of the limited resolution, these events could leak into the fit region from  $-0.12$  to  $0.30$  GeV. We determine the resulting systematic uncertainty by changing the fit region to  $-0.10$  to  $0.30$  GeV. The branching fraction changes by 1.8%.

The uncertainty arising from the chosen background description is determined by repeating the fit to determine

the signal yield with a second-order polynomial for the background. The number of signal events changes by 2.0%. A comparison between the data and Monte Carlo events shows that the mean of the  $\Delta E$  distribution in the data is shifted by  $-0.003$  GeV. We determine the resulting systematic uncertainty by shifting the signal region for the fit in the Monte Carlo  $\Delta E$  distribution by  $0.003$  GeV. This changes the efficiency to 8.60%, corresponding to a systematic uncertainty of 0.8%.

The uncertainty due to the treatment of the three-body phase space in the efficiency correction is estimated from the variation of the efficiency when performing further iterations of weighting. Table I shows a variation from 8.81% down to 8.77%. This corresponds to a systematic uncertainty of 0.5%.

A possible additional contribution to the statistical uncertainty coming from the efficiency determination, where we determined the weights based on data events in ranges of the two-body masses, is studied by performing the efficiency correction in ranges of  $m(\Lambda_c^+ K^-)$  only, with unweighted Monte Carlo events. The change in overall reconstruction efficiency is negligible compared to the statistical uncertainty.

Adding all contributions in quadrature we obtain a systematic uncertainty of 4.1%. The significance of the signal, including additive systematic uncertainties, is determined to be more than 7 standard deviations. This significance

includes systematic uncertainties from the  $\bar{B}^0 \rightarrow \Lambda_c^+ \bar{\Sigma}^0 K^-$  and  $B \rightarrow \Lambda_c^+ \bar{\Lambda} K^- \pi$  background as well as the  $\Delta E$  background description.

In summary, we observe the decay  $\bar{B}^0 \rightarrow \Lambda_c^+ \bar{\Lambda} K^-$  with a significance larger than 7 standard deviations and measure a branching fraction of

$$\mathcal{B}(\bar{B}^0 \rightarrow \Lambda_c^+ \bar{\Lambda} K^-) = (3.8 \pm 0.8_{\text{stat}} \pm 0.2_{\text{sys}} \pm 1.0_{\Lambda_c^+}) \times 10^{-5}. \quad (3)$$

The decay rate is not uniform over three-body phase space; rather, it is dominant at high  $\Lambda_c^+ K^-$  mass.

We are grateful for the excellent luminosity and machine conditions provided by our PEP-II colleagues and for the substantial dedicated effort from the computing organizations that support *BABAR*. The collaborating institutions thank SLAC for its support and kind hospitality. This work is supported by DOE and NSF (USA), NSERC (Canada), CEA and CNRS-IN2P3 (France), BMBF and DFG (Germany), INFN (Italy), FOM (The Netherlands), NFR (Norway), MES (Russia), MICINN (Spain), and STFC (United Kingdom). Individuals have received support from the Marie Curie EIF (European Union), the A.P. Sloan Foundation (USA), and the Binational Science Foundation (USA–Israel). The collaborating institutions thank SLAC for its support and kind hospitality.

- 
- [1] K. Nakamura *et al.* (Particle Data Group), *J. Phys. G* **37**, 075021 (2010).
  - [2] Throughout this paper, all decay modes represent that mode and its charge conjugate.
  - [3] D.J. Lange, *Nucl. Instrum. Methods Phys. Res., Sect. A* **462**, 152 (2001).
  - [4] S. Agostinelli *et al.* (GEANT4 Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **506**, 250 (2003).

- [5] B. Aubert *et al.* (*BABAR* Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **479**, 1 (2002).
- [6] B. Aubert *et al.* (*BABAR* Collaboration), *Phys. Rev. Lett.* **99**, 021603 (2007).
- [7] B. Aubert *et al.* (*BABAR* Collaboration), *Phys. Rev. D* **80**, 051105 (2009).